

For the use of Registered Veterinary Practitioner, Hospital or Laboratory

FIPROFORT® SPRAY

(Fipronil 0.25%w/v spray)

Composition:

Each ml of FIPROFORT® Spray contains:

Fipronil	0.25% w/v
Excipients	q.s.

Pharmacology:

Mechanism of action:

Fipronil is a new generation broad-spectrum ectoparasitocidal belonging to phenylpyrazole class. Fipronil disrupts the insect's central nervous system by blocking the passage of chloride ions through the GABA receptor and glutamate receptor (GluCl).

Fipronil acts by binding to an allosteric site of GABA receptors. Its neurotoxicity is selective, because the configuration of GABA receptors in mammals is different from insects.

Indications:

For effective treatment and prevention of all life stages of ectoparasites i.e. tick (including ticks responsible for tick fever), flea (flea allergy dermatitis) and lice in Dogs, Puppies, Cats and Kittens.

Application:

In order to wet the coat down to the skin it is recommended that the following application rates be used:

Short-haired animals (<1.5 cm)
A minimum of 3 ml/kg body mass =
7.5 mg of active material kg/body mass.

Long-haired animals (>1.5 cm)
A minimum of 6 ml/kg body mass =
15 mg of active material kg/body mass.

100 ml bottle

Each actuation delivers 0.5 ml spray volume
6 pumps actions per kg
e.g. for dogs under 12 kg
Weight 15 kg = 90 pump actions
Weight 30 kg = 180 pump actions

250 ml bottle

Each trigger application delivers 1 ml spray volume
3 pump actions per kg
e.g. for large dogs over 12 kg

Weight 15 kg = 45 pumps actions

Weight 30 kg = 90 pumps actions

Divide the animals body into five parts, right side, left side, underbelly, head to tail and face. An equal number of pumps to be administered in each of the part.

Directions for use:

TO PREVENT HARM TO YOU AND YOUR PET, READ ENTIRE LABEL AND ATTACHED DIRECTIONS BEFORE EACH USE.

Follow all directions and precautionary statements carefully. Use only on dogs and cats. Do not use on rabbits. Do not use on any other animals.

For effective treatment and prevention of all life stages of ectoparasites i.e. tick (including ticks responsible for tick fever), flea (flea allergy dermatitis) and chewing lice, apply FIPROFORT Spray to dogs, cats, puppies and kittens as follows:

For 100 ml and 250 ml

Hold the bottle in upright position. Ruffle the animal's coat while applying spray mist to its body.

Put a pair of disposable gloves.

Spray on the body of animal from a distance of 10-20 cm against the direction of hair in a well-ventilated room (if you are treating a dog, you may prefer to treat it outside).

Apply on entire body concentrating on affected area. Coat the spray all over to make sure the spray gets right down to the skin.

Allow animal to air dry. Do not towel dry.

Residual action:

Ticks: 1 month (3-5 week), Fleas: 2 months (1-3 months) this will depend on the parasite challenge in the animal's environment.

Precautions:

- Avoid spraying into eyes while spraying on face. To prevent spraying into eyes and to ensure proper coverage on the head in nervous animals, puppies and kittens spray Fiprofort onto your gloves and rub on face and other body parts.
- Do not allow animals to lick the spray.

- Do not shampoo for at least 2 days before and after Fiprofort treatment.
- Do not smoke, eat or drink during application.
- Wear gloves during spraying.
- Wash the hands after use.
- Spray in a well-ventilated area.
- Keep sprayed animals away from the source of heat until the animal gets dry.
- Do not spray directly on the area of damaged skin.

Safety:

FIPROFORT® Spray is safe to use in dogs and cats with no toxic effects when used in recommended way. FIPROFORT® Spray is not absorbed by skin and stay on for longer duration to provide effective protection. FIPROFORT® Spray is safe to use on cats and dogs 8 weeks of age and older, pregnant and lactating bitches.

Toxicity:

In case of accidental ingestion or excessive licking by your pet contact your veterinarian immediately for symptomatic treatment. Wash with plenty of fresh water immediately in case of contact with eyes or open wound.

Only for use by or under the control of a veterinarian.

Storage: Store below 30°C. Keep out of reach of children.

Container disposal:

Empty packages or unused products can be disposed at the waste disposal facility. Do not reuse this container.

Physical or Chemical Hazards: FLAMMABLE
Keep away from heat and open flame.

Presentation: Bottle of 100 ml and 250 ml

Last Update: July 2013



SAVA HEALTHCARE LIMITED

Plot No. 507-B to 512 GIDC Estate,
Wadhwan City – 363035, Surendranagar,
Gujarat, India

www.savavet.com