

For the use of Registered Veterinary Practitioner or a Hospital or a Laboratory only

OBLIXIN LA
(Enrofloxacin LA 10% Injection)

Ready To Use Antibiotic Injection for treatment of bacterial infection in animals.

Composition

Each ml contains:

Composition:

Each ml contains

Enrofloxacin IP- 100mg

Benzyl alcohol IP- 1.5%w/v

Long-Acting Vehicles- q.s.

Pharmacological Properties:

Oblixin-LA contains Enrofloxacin, which is a third-generation fluoroquinolone offering a broad spectrum of bactericidal activity against Gram-positive and/or Gram-negative bacteria and Mycoplasma species. Enrofloxacin acts by its specific interfering property of the DNA gyrase enzyme and thereby inhibits the metabolism of bacteria.

The formulation of Oblixin-LA offers the following advantages

- Excellent bioavailability
- Long-acting efficacy
- Rapid diffusion into all parts of the body
- Better tolerance and stability

The Ready to Use formulation allows avoiding the dilution before the injection and helps the ease of use. The Oblixin-LA formulation also allows reducing the number of injections in contrast to the traditional Enrofloxacin Injections. Only one injection of Oblixin-LA is usually sufficient for complete treatment. If required, it can be repeated after 72 hours

of first injection, and thus it helps in therapeutic compliance.

Dosage:

Ruminants & Pigs

One Injection of 3 ml / 40 kg of body weight or 7.5 mg / kg body weight. Repeat after 72 hours, if necessary.

Sheep & Goat

One Injection of 1.5 ml / 20 kg body weight or 7.5 mg / kg body weight. Repeat after 72 hours, if necessary. Or as directed by the Veterinarian.

ADMINISTRATION

For IV/IM/SC Use Only.

Administered dose should not exceed

*15 ml at a single site in cattle, buffalo & camels

*5 ml at a single site in sheep, goats & pigs

Presentation: 30 ml, & 100 ml glass vial

Indications:

Oblixin-LA is recommended for the treatment of

- Respiratory Diseases
- Digestive Diseases
- Other diseases caused by both Gram positive and/or Gram negative organisms of primary and secondary nature.

Oblixin-LA is also effective against the following microorganisms:

Mycoplasma spp., *E. coli*, *Salmonella* spp.
Pasteurella spp. *Mannheimia haemolytica*,

Bordetella spp., *Actinobacillus pleuropneumoniae*, *Enterobacteriae*, *Pyogenes* bacteria, *Streptococcus* spp. *Staphylococcus* spp.

Withdrawal period:

Milk: 7 days, **Meat:** 14 days

Carefully read the accompanying instructions before use.

**VETERINARY | NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**



SAVA HEALTHCARE LIMITED
Plot No. 507-B to 512, G.I.D.C. Estate,
Wadhwan City-363035, Surendranagar,
Gujarat. INDIA. | www.savavet.com